



The Central Role of the Radiology Nurse in Ensuring Safe Administration and Handling of Contrast Media: Systematic Review

Yasser Said Alzahrani⁽¹⁾, Zainab Abdullah Saleh Albishi⁽¹⁾, Majeb Mubark Aldosari⁽²⁾, Zahra Ahmed Aysh Alsmail⁽³⁾, Ahmed Ibrahim Ibrahim Jarrah⁽⁴⁾, Haya Saad Said Aldossari⁽⁵⁾, Abdullah Hamood Alazmy⁽⁶⁾, Ibtesam Ali Alwan⁽⁷⁾, Majed Kasab Shlman Alanazi⁽⁸⁾, Mody Nasser Alkhamiss⁽⁹⁾, Latifa Abdullah Al-Jouraid⁽⁹⁾, Abdullah Fahad Alntaifat⁽¹⁰⁾, Sawsan Hussain Mohammed Alawad⁽¹¹⁾

⁽¹⁾Al-Sulayyil General Hospital, Ministry of Health, Saudi Arabia

⁽²⁾Prince Sultan Health Centre, Ministry of Health, Saudi Arabia

⁽³⁾King Fahad Hospital-Aljaber Kidney Center, Ministry of Health, Saudi Arabia

⁽⁴⁾Al-Harth General Hospital, Ministry of Health, Saudi Arabia

⁽⁵⁾King Khalid Hospital In Al Kharj, Ministry of Health, Saudi Arabia

⁽⁶⁾Nafi General Hospital, Ministry of Health, Saudi Arabia

⁽⁷⁾Alsuwedi Primary Health Center, Ministry of Health, Saudi Arabia

⁽⁸⁾Rumah Hospital, Ministry of Health, Saudi Arabia

⁽⁹⁾Alhair Phc, Ministry of Health, Saudi Arabia

⁽¹⁰⁾Ministry of Health, Saudi Arabia

⁽¹¹⁾Maternity and Children Hospital Alahsa, Ministry of Health, Saudi Arabia

Abstract

Background: Contrast media administration is a pillar of current diagnostic and interventional radiology to maximize image quality and diagnostic information. It is associated with a variety of risks, from harmless side effects to life-threatening emergencies. Within the multimodal imaging team, the radiology nurse has evolved into a pivotal player, acting as the patient safety key throughout the periprocedural path.

Aim: This systematic review claims to synthesize available evidence and formulate guidelines to outline the radiology nurse's multifaceted role in the safe administration and management of contrast media. It claims to write about the underlying tasks that define this role and highlight the nurse's vital role in minimizing risk and improving patient outcomes.

Methods: A systematic integration and synthesis of evidence from a wide range of peer-reviewed literature, including professional guidelines from associations such as the American College of Radiology and the European Society of Urogenital Radiology, was conducted. The review systematically reviews the evidence-based practices that apply to each aspect of the role of the radiology nurse.

Results: The review details and summarizes the key responsibilities of the radiology nurse, which form a cohesive framework for maintaining patient safety. They include pre-procedure patient evaluation and screening, knowledge of pharmacology related to contrast medium administration, application of accurate administration methods, careful patient monitoring, and management of emergent adverse effects. Additionally, the results highlight the nurse's key roles of patient education, advocacy, and facilitating a culture of safety and interprofessional communication within the radiology department.

Conclusion: The radiology nurse is crucial to effective and safe administration of contrast media. Expertise from the initial risk assessment and prevention all the way to emergency handling by the radiology nurse significantly decreases the intrinsic risks of contrast administration. By guaranteeing safety measures are being obeyed and patient care is being considered as a whole, the radiology nurse is part of the successful patient outcomes in modern radiological practice.

Keywords: Radiology Nurse, Contrast Media, Patient Safety, Adverse Reactions, Contrast-Induced Nephropathy, Extravasation, Pre-Procedural Screening, Emergency Management, Interprofessional Collaboration, Guidelines..

1. Introduction

Diagnostic and interventional radiology has undergone a revolution in recent decades and has become an essential component of modern medicine. In the center of this change is intravascular contrast media (CM), or drug agents administered to patients to

enhance contrast between anatomical structures or pathological tissues on radiographic images (Brant & Helms, 2007). Application of CM, including iodinated contrast media (ICM) for computed tomography (CT) and fluoroscopy, and gadolinium-based contrast agents (GBCA) for magnetic resonance imaging

(MRI), has increased exponentially over the years, with millions of doses being administered yearly worldwide (Davenport et al., 2020).

Despite being safe in the majority of instances, the application of CM is a medical intervention with intrinsic risks. These range from mild, self-limited reactions to severe, life-threatening emergencies such as anaphylactoid reactions, contrast-induced nephropathy (CIN), or extensive tissue damage caused by extravasation (Khan et al., 2023; Behzadi et al., 2018). The patients presenting to radiology units are also increasingly complex, with more comorbidities such as renal impairment, diabetes, heart failure, and multiple allergies, all serving to enhance the burden of risk for CM injection (Wang et al., 2008).

To meet these needs, the radiology nurse professional role has shifted from the periphery of an assistant to a central imaging team member. The radiology nurse is an advanced knowledge and skill registered nurse in the practice of patient care in the radiology environment (Lehmann, 2019). Their expertise is at the core of directing the entire patient pathway, from pre-procedure assessment to post-procedure care, with an ongoing focus on risk minimization and patient safety.

This review aims to provide a comprehensive discussion of guidelines and evidence-based practice that define the position of the radiology nurse in the safe handling and administration of contrast media. Through the combination of literature today and professional standards, this review will sanctify the radiology nurse as a pre-eminent protector through whom the diagnostic benefits of contrast media are achieved at the lowest possible risk to the patient.

Pre-Procedural Patient Evaluation and Risk Stratification

The contrast media safe use is not a process beginning in the exam room but during the meticulous pre-procedural period, a phase extensively organized and executed by the radiology nurse. This is an important phase for risk identification and prevention, whereby formal and systematic assessment is the optimal method of evading adverse events (Werthman, 2019). The contribution of the nurse in this sector is three-fold, beginning with a thorough history-taking that serves as the foundation for patient safety. The most crucial is a comprehensive allergy history, with specific attention to any past allergic or anaphylactic reactions to medication, food, or airborne allergens. Perhaps most critical is a past reaction to the contrast medium itself, which is the single largest risk factor for a future reaction, increasing the risk by 3 to 5-fold, and this must be documented carefully regarding the type, severity, and management needed of the reaction (Brockow, 2021).

Simultaneously, the nurse must conduct a comprehensive renal function test to identify patients at risk of Contrast-Induced Nephropathy (CIN) or, in the case of using MRI, Nephrogenic Systemic Fibrosis

(NSF). This entails requesting a history of chronic kidney disease, diabetes mellitus, hypertension, heart failure, dehydration, and concomitant use of nephrotoxic drugs such as metformin and NSAIDs (Mehran & Nikolsky, 2006; Petrov et al., 2022). In accordance with established standards of the American College of Radiology (ACR) and European Society of Urogenital Radiology (ESUR), the nurse determines an estimated Glomerular Filtration Rate (eGFR) in each at-risk patient prior to the administration of iodinated contrast media (ACR, 2021; ESUR, 2022).

In addition to renal function and allergy, the question includes the cardiovascular and respiratory systems. These conditions are significant to identify because they have the potential to either exacerbate the contrast injection-induced physiological stress or mimic the presentation of a severe adverse reaction (de Santa María et al., 2023). Reviewing all of the medications present, including herbal supplements and over-the-counter medications, is also of great importance. The nurse is especially vigilant with beta-blockers, which will mask the tachycardic reaction typical of an anaphylactic reaction and render it more refractory to conventional therapy with epinephrine, and with metformin, which requires specific withholding and re-initiation strategies post-procedure to prevent the rare but serious complication of lactic acidosis (ACR, 2021).

This data collection is augmented by a focused physical and psychosocial assessment. The nurse evaluates locations for intravenous (IV) access for appropriateness, takes baseline vital signs to have a valuable point of reference during the procedure, and determines physical limitations. Furthermore, the nurse evaluates the patient's level of anxiety and knowledge regarding the impending procedure. This psychosocial assessment is clinically relevant, since anxious patients are more likely to experience vasovagal responses, whose symptoms, such as bradycardia and hypotension, may be mistaken for more severe side effects; timely communication and reassurance can greatly enhance patient compliance and satisfaction (Thim et al., 2022). The radiology nurse also has an irreplaceable role in informed consent. Though the radiologist has the final say regarding the medical benefits and risks, the nurse has a patient advocate role, ensuring the information is indeed understood, all questions are answered, and the consent is informed and voluntary, hence upholding the ethical integrity of the procedure (Doudenkova & Bélisle Pilon, 2016).

Summarizing this comprehensive review, the radiology nurse then stratifies the patient's overall risk and collaborates directly with the radiologist in implementing individualized measures of prevention. For those patients with a history of acute contrast media reaction, this involves coordinating and administering a premedication regimen involving corticosteroids and antihistamines, as briefly outlined

in ACR guidelines, typically administered 13 and 7 hours prior to the procedure (ACR, 2021). For those patients found to have renal impairment, the nurse coordinates and regularly administers pre-emptive hydration protocols, oral or IV, which is the most evidence-based intervention for avoiding Contrast-Induced Nephropathy (Kaliyaperumal et al., 2023; Murugan et al., 2023). In doing so, the radiology nurse takes raw patient data and translates it into a living and active safety plan, essentially setting the stage for a safe and effective imaging study. Figure 1 illustrates the radiology nurse's role in risk assessment and preparation.




Figure 1: Pre-Procedural Patient Safety Workflow Pharmacologic Understanding of Contrast Media

A deep and complex understanding of the pharmacologic characteristics of the various contrast media is a fundamental pillar of the radiology nurse's knowledge, directly informing each aspect of their clinical practice, ranging from decision-making and monitoring to rescue. It begins with an understanding of the contrasting classes of contrast media. Iodinated Contrast Media (ICM), used in computed tomography (CT), fluoroscopy, and angiography, are generally classified according to their osmolality relative to blood plasma. High-Osmolar Contrast Media (HOCM) are 5-8 times more osmolal than blood and are older compounds no longer widely used for intravascular administration since they are strongly associated with a higher risk of adverse reactions. Low-Osmolar Contrast Media (LOCM) with 2-3 times blood osmolality are the current practice standard, providing significantly improved safety against nephrotoxicity and allergic-like responses. Iso-Osmolar Contrast Media (IOCM) with matching osmolality to blood are reserved for specific high-risk cases, such as cardiac and renal angiography in

vulnerable patients (Davenport et al., 2020; Zamora & Castillo, 2017). For Magnetic Resonance Imaging (MRI), Gadolinium-Based Contrast Agents (GBCA) are used, and the nurse should recognize the significant distinction between linear and macrocyclic agents. Macrocyclic agents are more stable chemically, which is concerning a significantly reduced risk of causing Nephrogenic Systemic Fibrosis (NSF) in patients with extremely impaired renal function, which determines the selection of agent in such at-risk patients (ESUR, 2022).

The radiology nurse is aware beyond simple categorization to the physiologic effects and inherent toxicity of these drugs. When given intravenously, contrast medium creates a massive osmotic load, with predictable physiological consequences such as peripheral vasodilation, transient increase in cardiac output, and sensation of heat or flushing; knowledge of the expected effects is the basis for the nurse to be able to distinguish them confidently from the early manifestations of an abnormal adverse effect (Aggarwal, 2023). These are the primary mechanisms of toxicity: chemotoxicity, the direct irritant effect of the contrast molecule itself on tissues and organs; osmotoxicity, relating to the damage which hyperosmolar solutions can inflict on endothelial cells and red cells; and viscosity, a physical property that governs flow dynamics. Higher viscosity agents require the use of larger-bore IV catheters and create higher injection pressures and, therefore, contribute directly to the mechanical risk of extravasation (Lightfoot et al., 2009). All this pharmacologic knowledge base ultimately allows the radiology nurse to be able to anticipate possible complications, make rational decisions about equipment selection, and provide patients and peers with rational, evidence-based justifications.

Safe Preparation, Handling, and Administration Techniques

The actual procedure of producing and administering contrast media is a high-risk process that needs precise adherence to aseptic technique and stringent safety measures, where the radiology nurse stands as the key professional who guarantees that standards are always maintained. Preparation is governed by meticulous attention to detail. The nurse must ensure the appropriate contrast agent, its concentration, volume, and expiration date against the physician's order, a necessary step in averting medication errors (Hecht et al., 2006). This is followed by the use of contrast media from sterile syringes and strict aseptic technique to prevent microbial contamination. Besides, the nurse should also possess a good understanding of the incompatibility of drugs so they recognize that contrast media should never be used in conjunction with other drugs in the same syringe or IV line since precipitation and crystallization are extremely probable (SRNA, 2022).

Vascular access management is likely one of the most personal and significant responsibilities of the radiology nurse in this context. Proper IV access is paramount to the success of the study and to complication avoidance. The nurse performs a competent venous system assessment, selecting a vein based on size, integrity, and location. The most favorable sites are the antecubital fossa or large forearm vein and offer consistent blood flow and are less likely to shift. Conversely, dorsal hand veins or veins around mobile joints are considered inferior due to their smaller size and higher chances of related contrast extravasation (Ding et al., 2018). The nurse then selects an appropriately sized IV catheter—typically an 18- or 20-gauge for power-injected CT studies to permit high flow rates—secures it firmly to the patient's skin, and gives a final definitive saline flush to test patency, observing for resistance, swelling, or patient distress, an extremely important final test before going on.

The actual process of delivering contrast media is a technical procedure, increasingly mastered with automated power injectors. The radiology nurse must be completely trained on the use, programming, and safety features of such machines. This involves accurately setting the parameters of injection—flow rate (in mL/sec), total volume (in mL), and pressure

limit—tailored to the appropriate protocol and patient vasculature. One of the non-negotiable safety precautions is that all connections within the laboratory between the IV catheter, syringe, and tubing are Luer-lock fittings, which are mandatory to prevent catastrophic disconnection from the high pressures generated from the injector (Wang et al., 2017). One aspect of the administration procedure is foresight in talking to the patient. The nurse administers the injection just before and then prepares the patient for the very normal sensations that they will be experiencing, which are a flush of heat, a metallic taste in the mouth, or a fleeting sensation of having wet themselves. This basic act of teaching is an extremely effective risk-reduction measure, since it reduces patients' anxiety levels substantially and prevents the abrupt, startled response that is a common cause of IV dislodgement and contrast extravasation (Werthman et al., 2021). Through this combination of technical competence, planning, and compassionate communication, the radiology nurse renders the physical administration of contrast media as secure as possible (Table 1). Figure 2 summarizes the essential pharmacologic principles guiding the safe use of contrast media in radiology

Table 1: Key Steps for Safe Contrast Media Administration by the Radiology Nurse

Phase	Key Nursing Action	Rationale
Pre- Procedural	Perform comprehensive patient assessment (allergies, renal function, medications).	To identify risk factors and stratify patient for adverse events.
	Verify informed consent is obtained and understood.	To uphold ethical standards and ensure patient autonomy.
	Obtain and document baseline vital signs.	To provide a comparison for intra- and post-procedural monitoring.
	Establish patent IV access with appropriate gauge catheter.	To ensure reliable venous access for injection and minimize extravasation risk.
Intra- Procedural	Confirm correct patient, contrast agent, and dose using two patient identifiers.	To prevent medication errors.
	Use Luer-lock connections on all syringe and tubing attachments.	To prevent disconnection under high pressure from power injectors.
	Educate the patient on expected sensations during injection.	To minimize anxiety and prevent motion artifact or sudden movement.
	Maintain direct visual or verbal contact with the patient during injection.	To allow for immediate recognition of adverse reactions or extravasation.
Post- Procedural	Monitor patient in the department for a period (e.g., 15-30 minutes).	To identify delayed adverse reactions.
	Perform a post-injection site assessment.	To check for signs of delayed extravasation or phlebitis.
	Provide post-procedure discharge instructions, both verbal and written.	To empower the patient with knowledge of signs/symptoms to watch for at home.
	Ensure accurate and timely documentation of the entire process.	To create a legal record and facilitate communication with the care team.

Management and Monitoring of the Patient During the Procedure

Active and continuous patient monitoring while and immediately after contrast media administration is a primary and non-delegable role of

the radiology nurse. Careful and vigilant monitoring is the most crucial safety net to allow detection of any deviation from the expected physiological response at the earliest moment. The nurse assessment is an active and thorough process that seeks to catch subtle

indicators, which might be the beginning of a negative reaction. This includes systematic examination for cutaneous findings such as urticaria, flushing, or pruritus; respiratory findings such as a new cough, sneezing, nasal congestion, hoarseness, wheezing, or the sinister finding of stridor indicative of upper airway compromise; and cardiovascular changes such as tachycardia, bradycardia (which is characteristic of a vasovagal response), hypotension, or hypertension. The nurse also monitors for neurological manifestations of heightened anxiety, dizziness, or syncope, subjective complaints by the patient, including nausea, vomiting, or a feeling of intense "impending doom," which may be an intense predictive sign of a severe reaction (Bush, 1990). Presence of the nurse and frequent, competent monitoring are the important first line of defense in the radiology suite, allowing for prompt intervention before a mild, local reaction turns into a severe, systemic, or life-threatening one.




Figure 2. The essential pharmacologic principles guiding the safe use of contrast media in radiology.

Recognition and Management of Acute Adverse Reactions

Despite the strictest pre-procedure prevention practices, unpredictable contrast media adverse reactions can and do occur. The radiology nurse must, therefore, possess high levels of competency in the rapid assessment, grading, and initial management of such incidents. For seamless and effective reaction, the entire radiology staff, with the nurse as a central figure, would require training in emergency procedures and participate in regular simulated drills to remain current (SRNA, 2022).

Adverse reactions are typically classified based on their timing and pathophysiologic

mechanism. They are acute within one hour of injection and can be either allergic-like (anaphylactoid), which are not IgE-mediated but clinically resemble them, or non-allergic (chemotoxic), caused by the direct toxic or osmotic action of the agent. It is of utmost significance to note that from a clinical management perspective, it is often not feasible to discriminate between a true IgE-mediated anaphylactic reaction and an anaphylactoid reaction; therefore, both are handled in the same manner with the same level of urgency and management intervention (Brockow, 2003). Reactions occur between one hour to seven days post-injection. These are typically milder and self-limiting, commonly appearing as cutaneous reactions like a mild rash but also including nausea, headache, and flu-like syndrome (ACR, 2021).

During an acute reaction, the radiology nurse is always the first to respond, in charge of initiating life-saving actions while simultaneously calling the radiologist and initiating other medical intervention. Its management philosophy is firmly rooted in the ABCs (Airway, Breathing, Circulation) of advanced life support. For mild reactions, such as minor urticaria or pruritus, management may consist of simple observation or administration of an oral or intramuscular antihistamine such as diphenhydramine with close observation by the nurse for deterioration (Lightfoot et al., 2009). Moderate reactions, with diffuse urticaria, mild bronchospasm, or facial edema, necessitate a more aggressive and systematic approach. The nurse administers supplemental oxygen at once, checks IV access as patent and adequate, and administers medications per protocol, which may be antihistamines and corticosteroids, while scrupulously observing and documenting vital signs (Wahl et al., 2023).

In the event of an extreme response—such as laryngeal edema, intense bronchospasm, hypotension leading to shock, or cardiac arrest—the nurse's intervention becomes absolutely imperative. These are full-scale medical emergencies requiring drastic and immediate action. The nurse's responsibility is the management of the airway by way of high-flow oxygen administration and prepping for advanced airway management in case laryngeal edema is a suspicion; provision of assistance in breathing by way of beta-agonist bronchodilator such as albuterol through the nebulizer for bronchospasm, with subcutaneous or intramuscular epinephrine being the first agent for severe cases; and maintenance of the circulation through positioning of the patient in Trendelenburg position, rapid intravenous bolus, and preparation to administer epinephrine intramuscularly or as an intravenous infusion in case of refractory hypotension (Hsieh et al., 2022; ACR, 2021). To assist in this emergency response, the radiology unit must have a well-stocked, properly labeled, and regularly inspected "contrast reaction" or "crash" cart. The

radiology nurse is likely also to be the sharer of this precious asset, ensuring all drugs and equipment are within their expiration dates and immediately available for use at the time of a crisis (Table 2).

Table 2: Management of Acute Adverse Reactions to Contrast Media: A Nursing Guide

Reaction Severity	Signs and Symptoms	Initial Nursing Actions & Medications
Mild	Limited urticaria, pruritus, minor flushing, rhinorrhea.	1. Reassure the patient. 2. Monitor for progression. 3. Consider diphenhydramine 25-50 mg PO/IM/IV.
Moderate	Diffuse urticaria, mild bronchospasm (wheezing), facial edema, tachycardia, hypertension.	1. Maintain patient airway. Administer O ₂ 6-10 L/min via mask. 2. Establish/ensure IV access. 3. Bronchospasm: Albuterol 2.5 mg via nebulizer. 4. Urticaria/Edema: Diphenhydramine 25-50 mg IM/IV. 5. Consider: Ranitidine 50 mg IV or Methylprednisolone 125 mg IV.
Severe (Anaphylactoid)	Laryngeal edema (stridor), profound bronchospasm, hypotension/shock (SBP <90mmHg), arrhythmias, cardiac arrest.	1. CALL A CODE/CRASH TEAM. 2. Airway/Breathing: High-flow O ₂ (100%). Prepare for intubation. Laryngeal Edema/Bronchospasm/Shock: Epinephrine 1:1,000, 0.3-0.5 mg IM in the mid-antrolateral thigh. Repeat every 5-15 mins as needed. 3. Circulation: Place in supine/Trendelenburg. Rapid IV fluid bolus (0.9% NaCl). For refractory hypotension, prepare epinephrine IV infusion. 4. Cardiac Arrest: Initiate CPR and follow ACLS protocols.
Vasovagal Reaction	Bradycardia, hypotension, diaphoresis, nausea, syncope.	1. Place patient in Trendelenburg position. 2. Administer O ₂ . 3. Administer rapid IV fluid bolus. 4. If bradycardia persists: Atropine 0.6-1.0 mg IV, repeat to a max of 3 mg.

Prevention and Prevention of Non-Acute Complications

Along with acute allergic-like reactions, early treatment, the radiology nurse plays a pivotal and active role in the prevention and control of other severe, non-acute complications involving contrast media. Possibly the most serious among them is Contrast-Induced Nephropathy (CIN), an acute renal injury that typically develops between 48-72 hours following intravascular iodinated contrast. The radiology nurse is most vital in preventing it, first by identifying at-risk patients during the pre-procedure evaluation, and then by triggering evidence-based preventive measures. The most critical of these are coordination and intravenous hydration with 0.9% sodium chloride solution, which is still the most effective preventive measure. This is generally taken in the form of a standard protocol of 1 mL/kg/hr over 6-12 hours prior to and after the procedure, or a rapid bolus of 3-5 mL/kg over one hour taken immediately before the study (Weisbord & Palevsky, 2020; Weisbord, 2018). The nurse also facilitates medication management by counseling withholding nephrotoxic agents like NSAIDs on a temporary basis and coordinating with the referring doctor for metformin, with particular withholding and reinitiating protocols required in patients with compromised renal function to prevent the risk of lactic acidosis (ACR, 2021).

Another common complication in which the radiology nurse has an important role is extravasation, that is, unintentional leakage of contrast medium into surrounding soft tissues of the vein. While generally benign, extravasation can lead to severe pain, swelling, and, in extreme cases involving large volumes of contrast medium, compartment syndrome or tissue necrosis. The nurse's function is varied, ranging from prevention through judicious IV site selection and patency checking, to rapid detection through diligent patient and injection site monitoring. If extravasation occurs, the nurse adheres to standard management, which includes stopping the injection immediately, elevating the affected extremity to promote reabsorption of the fluid, and applying a warm or cold compress—institutional guidelines vary, but warm compresses are generally employed in the case of low-osmolar contrast media to promote vasodilation and distribution (Roditi et al., 2022). Documentation of the event, including the estimated volume, specific location, and patient symptomatology, is required. For large-volume extravasations (usually >100 mL) or for any signs of neurovascular compromise, such as altered sensation, diminished pulse, or skin blisters, the nurse will get an urgent consultation with a plastic surgeon (ACR, 2021).

The radiology nurse is also a vital guardian against Nephrogenic Systemic Fibrosis (NSF), which is an uncommon but severe and possibly fatal fibrosing disease associated with particular gadolinium-based contrast media (GBCA) in individuals with severe renal impairment. The nurse prevents it through careful adherence to renal function screening prior to any GBCA administration and through encouragement of the use of the lowest dose of the most stable macrocyclic agent in such a patient population, a practice also advocated by international guidelines (ESUR, 2022).

Patient Education and Advocacy

Practice of the radiology nurse reaches deeply into the realm of patient education and advocacy, a steady and patient-centered pillar within the periprocedural process. Patient education is a process, not an event. In the pre-procedure time, the nurse provides appropriate information, including the reason for the test, demystifying the sensations of contrast injection, and reinforcing the absolutely essential necessity of the disclosure of a full and accurate medical history. Following the procedure, the nurse provides clear, concise, and often written post-procedure discharge instructions. Education post-procedure is required for ensuring patient safety upon discharge from the department; this includes teaching patients to take high amounts of fluid to enhance renal excretion of the contrast medium and explaining clearly when they should seek medical attention—such as by contacting their general practitioner or reporting back to the emergency department in the case of the onset of a delayed reaction (e.g., protracted rash, fever) or more dramatic symptoms at the injection site (e.g., pain, redness, blistering) (Park et al., 2023).

Parallel to education, the nurse is also an ongoing patient advocate, assuring that the patient's voice is heard and his or her rights and safety are the main priority. Such advocacy can take many forms, ranging from appealing a procedural order should a patient's risk factors appear unacceptably high, to delaying a procedure until there is absolute and genuine comprehension of the process of informed consent on the part of the patient. During emergencies, the nurse may be compelled to speak out assertively in order to get the appropriate treatment on time (Beckett et al., 2015). By this dual role as educator and advocate, the radiology nurse empowers patients and safeguards their health amidst the high-stakes setting of the radiology department.

Creating a Culture of Safety and Interprofessional Collaboration

Lastly, safety in the radiology suite is a responsibility that is shared, not individual, and the radiology nurse is a key supporter of a robust culture of safety and interprofessional collaboration. This culture is built on some key practices. Firstly, meticulous documentation of the entire patient

encounter from initial assessment and consent to precise parameters of contrast administration and response is an absolute necessity. This document is not just a legal and professional obligation but also a necessary tool for communication with the healthcare team and for quality improvement (SRNA, 2022).

Second, radiology nurses are engaged in Quality Improvement (QI) processes. They participate in tracking and assessing adverse event data, i.e., rate of reaction and occurrence of extravasations, in order to identify trends and system weaknesses. They also contribute to the review of emergency responses through debriefing and simulation exercises, all in the aim of identifying areas of improvement and ironing out clinical procedures (Tourgeman-Bashkin et al., 2013). Finally, the role of the nurse is essentially one of collaboration. They work seamlessly alongside radiologists, radiographers, and ancillary personnel, where respect and clear communication form the bases of both an efficient workflow and, most importantly, patient safety. Typically acting as the communication bridge between the patient, radiology department, and referring physician, the radiology nurse facilitates continuity of care and ensures that information pertinent to the case is properly conveyed, further solidifying their position as an integral integrator on the modern imaging team (Wang et al., 2017).

Conclusion

Contrast medium delivery, though mundane, is a highly advanced pharmacological treatment with great potential for patient harm. This article has critically spelled out the immense and critical role of the radiology nurse in avoiding these risks and guaranteeing of patient safety. From the first pre-procedure evaluation to high-stakes acute care of emergencies, the radiology nurse's specialized knowledge, clinical acumen, and vigilant eye are beyond price.

The radiology nurse is a risk-stratifier, educator, technical specialist, astute observer, emergency care provider, patient advocate, and safety champion. With an integrative evidence-based practice and professional guidelines model of care, their specialty brings the diagnostic benefits of contrast-enhanced imaging together with the highest possible degree of safety. With ongoing technological improvements in imaging and progressively more complicated patient populations, the role of the radiology nurse can only be more important. Continued investment in education, professional advancement encouragement, and official recognition of their unique contributions will be fundamental to pursuing further excellence and safety in radiological care.

References

1. Aggarwal, B. (2023). *Textbook of Radiology And Imaging, Volume 1-E-Book*. Elsevier Health Sciences.

2. American College of Radiology. (2021). *ACR manual on contrast media*. ACR Committee on Drugs and Contrast Media. <https://www.acr.org/Clinical-Resources/Clinical-Tools-and-Reference/Contrast-Manual>
3. Beckett, K. R., Moriarity, A. K., & Langer, J. M. (2015). Safe use of contrast media: what the radiologist needs to know. *Radiographics*, 35(6), 1738-1750. <https://doi.org/10.1148/rg.2015150033>
4. Behzadi, A. H., Zhao, Y., Farooq, Z., & Prince, M. R. (2018). Immediate allergic reactions to gadolinium-based contrast agents: a systematic review and meta-analysis. *Radiology*, 286(2), 471-482. <https://doi.org/10.1148/radiol.2017162740>
5. Brant, W. E., & Helms, C. A. (Eds.). (2007). *Fundamentals of diagnostic radiology*.
6. Brockow, K. (2005). Contrast media hypersensitivity—scope of the problem. *Toxicology*, 209(2), 189-192. <https://doi.org/10.1016/j.tox.2004.12.032>
7. Brockow, K. (2021). Diagnosing and managing patients with reactions to radiocontrast media. *Current Treatment Options in Allergy*, 8(3), 210-221. <https://doi.org/10.1007/s40521-021-00287-x>
8. Bush, W. H. (1990). Treatment of systemic reactions to contrast media. *Urology*, 35(2), 145-150. [https://doi.org/10.1016/0090-4295\(90\)80064-T](https://doi.org/10.1016/0090-4295(90)80064-T)
9. Davenport, M. S., Perazella, M. A., Yee, J., Dillman, J. R., Fine, D., McDonald, R. J., ... & Weinreb, J. C. (2020). Use of intravenous iodinated contrast media in patients with kidney disease: consensus statements from the American College of Radiology and the National Kidney Foundation. *Radiology*, 294(3), 660-668. <https://doi.org/10.1148/radiol.2019192094>
10. de Santa María, R. S., Labella, M., Bogas, G., Doña, I., & Torres, M. J. (2023). Hypersensitivity to gadolinium-based contrast. *Current opinion in allergy and clinical immunology*, 23(4), 300-306. DOI: 10.1097/ACI.0000000000000911
11. Ding, S., Meystre, N. R., Campeanu, C., & Gullo, G. (2018). Contrast media extravasations in patients undergoing computerized tomography scanning: a systematic review and meta-analysis of risk factors and interventions. *JBI Evidence Synthesis*, 16(1), 87-116. DOI: 10.11124/JBISIRIR-2017-003348
12. Doudenкова, V., & Bélisle Pipon, J. C. (2016, March). Duty to inform and informed consent in diagnostic radiology: how ethics and law can better guide practice. In *Hec forum* (Vol. 28, No. 1, pp. 75-94). Dordrecht: Springer Netherlands. <https://doi.org/10.1007/s10730-015-9275-7>
13. European Society of Urogenital Radiology. (2022). *ESUR contrast media guidelines, version 11.0*. <https://www.esur.org/esur-guidelines-on-contrast-agents/>
14. Hecht, E. M., Holland, A. E., Israel, G. M., Hahn, W. Y., Kim, D. C., West, A. B., ... & Krinsky, G. A. (2006). Hepatocellular carcinoma in the cirrhotic liver: gadolinium-enhanced 3D T1-weighted MR imaging as a stand-alone sequence for diagnosis. *Radiology*, 239(2), 438-447. <https://doi.org/10.1148/radiol.2392050551>
15. Hsieh, C., Wu, S. C., Kosik, R. O., Huang, Y. C., & Chan, W. P. (2022). Pharmacological prevention of hypersensitivity reactions caused by iodinated contrast media: a systematic review and meta-analysis. *Diagnostics*, 12(7), 1673. <https://doi.org/10.3390/diagnostics12071673>
16. Kaliyaperumal, Y., Sivadasan, S., & Aiyalu, R. (2023). Contrast-induced nephropathy: an overview. *Dr. Sulaiman Al Habib Medical Journal*, 5(4), 118-127. DOI: 10.1007/s44229-023-00040-3
17. Khan, F., Abbas, F., Hilal, K., Samad, M., Wahid, G., Ali, I., ... & Saeed, M. A. (2023). Knowledge assessment of radiologists, radiology residents, and radiographers regarding contrast materials and management of adverse drug reactions occurring due to contrast materials: a cross-sectional study. *Annals of Medicine and Surgery*, 85(7), 3347-3352. DOI: 10.1097/MS9.0000000000000465
18. Lehmann, S. L. (2019). Role of the Clinical Nurse Specialist in Radiology. In *Advanced Practice and Leadership in Radiology Nursing* (pp. 15-24). Cham: Springer International Publishing. https://doi.org/10.1007/978-3-030-32679-1_2
19. Lightfoot, C. B., Abraham, R. J., Mammen, T., Abdoell, M., Kapur, S., & Abraham, R. J. (2009). Survey of radiologists' knowledge regarding the management of severe contrast material-induced allergic reactions. *Radiology*, 251(3), 691-696. <https://doi.org/10.1148/radiol.2513081651>
20. Mehran, R., & Nikolsky, E. (2006). Contrast-induced nephropathy: definition, epidemiology, and patients at risk. *Kidney international*, 69, S11-S15. <https://doi.org/10.1038/sj.ki.5000368>
21. Murugan, R., Boudreaux-Kelly, M. Y., Kellum, J. A., Palevsky, P. M., & Weisbord, S. (2023). Contrast-associated acute kidney injury and cardiovascular events: a secondary analysis of the PRESERVE cohort. *Clinical*

- Kidney Journal*, 16(12), 2626-2638.
<https://doi.org/10.1093/ckj/sfad214>
22. Park, H. J., Shin, K., You, M. W., Kyung, S. G., Kim, S. Y., Park, S. H., ... & Kim, H. J. (2023). Deep learning-based detection of solid and cystic pancreatic neoplasms at contrast-enhanced CT. *Radiology*, 306(1), 140-149.
<https://doi.org/10.1148/radiol.220171>
 23. Petrov, V. I., Kudasheva, A. A., & Frolov, D. V. (2022). Contrast-induced nephropathy: prevalence, diagnosis, prevention and treatment. *Journal of Volgograd State Medical University*, 19(2), 7-18.
<https://doi.org/10.19163/1994-9480-2022-19-2-7-18>
 24. Roditi, G., Khan, N., van Der Molen, A. J., Bellin, M. F., Bertolotto, M., Brismar, T., ... & Clément, O. (2022). Intravenous contrast medium extravasation: systematic review and updated ESUR Contrast Media Safety Committee Guidelines. *European Radiology*, 32(5), 3056-3066.
<https://doi.org/10.1007/s00330-021-08433-4>
 25. Thim, S., Henriksen, T. B., Laursen, H., Schram, A. L., Paltved, C., & Lindhard, M. S. (2022). Simulation-based emergency team training in pediatrics: a systematic review. *Pediatrics*, 149(4), e2021054305.
<https://doi.org/10.1542/peds.2021-054305>
 26. Tourgeman-Bashkin, O., Shinar, D., Donchin, Y., Zmora, E., Velleman, N., & Libson, E. (2013). Radiology department, human factors and organizational perspectives: using action research to improve patient safety. *Israel Journal of health policy research*, 2(1), 40.
<https://doi.org/10.1186/2045-4015-2-40>
 27. Wahl, A. D., Saini, N., Trinh, K., Patel, S., Lobner, K., Huang, J., & Deng, F. (2023). Simulation training in the management of adverse contrast reactions: A systematic review. *Journal of the American College of Radiology*, 20(11), 1110-1120.
<https://doi.org/10.1016/j.jacr.2023.07.013>
 28. Wang, C. L., Chinnugounder, S., Hippe, D. S., Zaidi, S., O'Malley, R. B., Bhargava, P., & Bush, W. H. (2017). Comparative effectiveness of hands-on versus computer simulation-based training for contrast media reactions and teamwork skills. *Journal of the American College of Radiology*, 14(1), 103-110.
<https://doi.org/10.1016/j.jacr.2016.07.013>
 29. Wang, C. L., Cohan, R. H., Ellis, J. H., Caoili, E. M., Wang, G., & Francis, I. R. (2008). Frequency, outcome, and appropriateness of treatment of nonionic iodinated contrast media reactions. *American Journal of Roentgenology*, 191(2), 409-415.
<https://doi.org/10.2214/AJR.07.3421>
 30. Weisbord, S. D. (2018). Contrast-associated acute kidney injury. *Core Concepts in Acute Kidney Injury*, 95-112.
https://doi.org/10.1007/978-1-4939-8628-6_6
 31. Weisbord, S. D., Palevsky, P. M., Kaufman, J. S., Wu, H., Androsenko, M., Ferguson, R. E., ... & Gallagher, M. (2020). Contrast-associated acute kidney injury and serious adverse outcomes following angiography. *Journal of the American College of Cardiology*, 75(11), 1311-1320.
<https://doi.org/10.1016/j.jacc.2020.01.023>
 32. Werthman, J. A. (2019). An integrative review of outcomes and radiology nursing. *Journal of Radiology Nursing*, 38(1), 47-52.
<https://doi.org/10.1016/j.jradnu.2018.10.004>
 33. Werthman, J. A., Maxwell, C. A., Dietrich, M. S., Jordan, L. M., & Minnick, A. F. (2021). Moderate sedation education for nurses in interventional radiology to promote patient safety: results of a national survey. *Journal of Radiology Nursing*, 40(1), 49-55.
<https://doi.org/10.1016/j.jradnu.2020.10.007>
 34. Zamora, C. A., & Castillo, M. (2017). Historical perspective of imaging contrast agents. *Magnetic Resonance Imaging Clinics*, 25(4), 685-696.
<https://doi.org/10.1016/j.mric.2017.06.002>