Saudi Journal of Medicine and Public Health

https://saudijmph.com/index.php/pub

https://doi.org/10.64483/202412833

Interprofessional Strategies for Enhancing Patient Safety and Reducing Complications in Sedation and Anesthesia for Dental and Head-and-Neck Procedures: Roles of Laboratory Monitoring, Radiological Technology, Nursing, Physiotherapy, Dentistry, and Health Administration Systems

Yahya Ali Mugaddam Khabti[(1)] , Hind Abdulrahman Ahmed Al-Atiq[(2)] , Faisal Nawar Al-Harthi[(3)] , Nawal Mubarak Al-Khudran[(4)] , Yazeed Ali AlSanie[(5)] , Muteb Turki Saad Alotaibe[(6)] , Abdulrahman Abdullah A Alqahtani[(7)] , Abkar Ali Howash[(8)] , Eid Mohammed Ahmed Alnami[(9)] , Nouf Hassan Salman Al-Anazi[(10)]

(1)Sharourah General Hospital, Sharourah, Najran, Health Holding Company, Ministry of health, Saudi Arabia
(2)Ministry of Health, Saudi Arabia
(3)Riyadh Dental Complex, Ministry of Health, Riyadh First Health Cluster, Saudi Arabia
(4)Kood Asafyah Primary Health Care Center (PHCC), Ministry of health, Saudi Arabia
(5)Riyadh North dental center, Riyadh second health care cluster, Ministry of health, Saudi Arabia
(6)Almuzahmiah General Hospital, Ministry of Health, Saudi Arabia
(7)Khamis Mushayt Maternity and Children Hospital, Asir region, Ministry of health, Saudi Arabia
(8)Ministry of Health, Jazan Health Cluster, Tuberculosis Center, Saudi Arabia
(9)Ministry of Health, Al-Aidabi Hospital, Jazan, Sabya, Al-Hussaini, Saudi Arabia
(10)King Khalid Hospital, Ministry of Health, Al-Kharj, Saudi Arabia

Abstract

Sedation and anesthesia for dental and head-and-neck procedures take place in shared, anatomically constrained airways, frequently outside conventional operating rooms, and involve pediatric, geriatric, and oncologic patients who carry elevated risk. Complications such as hypoxemia, airway obstruction, aspiration, bleeding, delayed swallowing recovery, and communication failures are often preventable. This narrative review synthesizes evidence published between 2017 and 2025 on interprofessional strategies that strengthen safety, focusing on laboratory monitoring, radiological technology, nursing, physiotherapy, dentistry, and health administration systems. The literature indicates that structured teamwork, standardized monitoring including capnography, risk-stratified preoperative testing, imaging-informed airway planning, early-warning surveillance, prehabilitation and early rehabilitation, and governance that supports training, staffing, and incident learning are the most consistent contributors to risk reduction. An integrated model and implementation priorities are proposed, together with research gaps in outcome measurement, equity, and digital integration.

Keywords: interprofessional collaboration; patient safety; procedural sedation; dental anesthesia; head and neck surgery.

Introduction

Patient safety in anesthesia has improved substantially over recent decades, yet harm associated with health care remains a major global burden. A systematic review and meta-analysis estimated that roughly one in twenty patients experiences preventable harm across care settings, with a considerable proportion of this harm being severe or contributing to death (Panagioti et al., 2019). The World Health Organization has responded with a global action plan that frames safety as a system property requiring policy, culture, and cross-disciplinary engagement rather than the vigilance of individual clinicians alone (World Health Organization [WHO], 2021). The economic case is equally compelling, because avoidable harm consumes a large share of health expenditure and erodes institutional capacity (Slawomirski & Klazinga, 2022). Reflecting on two decades of progress since the landmark report on medical error, Bates and Singh (2018) argued that safety gains are most durable when organizations combine reliable processes, measurement, and teamwork.

Dental and head-and-neck procedures present a distinctive safety profile. The surgeon, dentist, or proceduralist and the anesthesia provider compete for access to the same airway, so that mouth, jaw, pharynx, larynx, and neck may simultaneously be the operative field and the route of oxygenation. Pathology, prior radiotherapy, trismus, tumor distortion, and fragile dentition complicate laryngoscopy and ventilation. Many dental interventions are delivered in ambulatory or office-based environments where emergency resources, laboratory support, and imaging are less immediately available than in hospital theatres. At the same time,

*Corresponding author yahyaali@gmail.com (Yahya Ali Mugaddam Khabti).
Receive Date: 22 December 2024, Revise Date: 30 December 2024, Accept Date: 31 December 2024

children requiring dental rehabilitation under sedation or general anesthesia and older adults with polypharmacy and multimorbidity represent populations with narrow physiological reserve (Coté & Wilson, 2019; Ashley et al., 2018).

The closed-claims and consensus literature repeatedly identifies airway-related events, inadequate monitoring, and failures of communication as recurring contributors to serious morbidity (Joffe et al., 2019; Cook & Harrop-Griffiths, 2019). These are not purely technical failures. They arise at the interfaces between professions: when a laboratory abnormality is not escalated, when imaging findings that predict a difficult airway do not reach the anesthesia team, when a nurse's observation of subtle deterioration is not acted upon, or when postoperative rehabilitation needs are not anticipated. A review of interprofessional collaboration confirms that structured collaborative interventions can improve professional practice and, in some contexts, patient outcomes, although effects depend on the quality of implementation (Reeves et al., 2017). Meta-analytic evidence also shows that better teamwork is associated with better clinical performance and fewer adverse events (Schmutz et al., 2019).

Much of the existing literature, however, examines professions in isolation. Guidelines for sedation, airway management, imaging, nursing surveillance, rehabilitation, and organizational governance are produced by separate communities, and few syntheses show how these contributions converge in a single patient pathway. The aim of this narrative review is therefore to integrate recent evidence (2017–2025) on how laboratory monitoring, radiological technology, nursing, physiotherapy, dentistry, and health administration systems can jointly reduce complications in sedation and anesthesia for dental and head-and-neck procedures, and to propose a practical framework for implementation.

2. Review Approach

This study follows a narrative review design, which is appropriate when the objective is conceptual integration across heterogeneous literatures rather than the pooling of effect sizes. Searches were conceptualized across major biomedical databases for publications from 2017 to 2025 using combinations of terms relating to procedural sedation, dental anesthesia, head-and-neck surgery, airway management, patient safety, interprofessional collaboration, laboratory testing, radiology, nursing, physiotherapy, and health system governance. Priority was given to clinical practice guidelines, consensus statements, systematic reviews, meta-analyses, large observational studies, and authoritative policy documents. Seminal older work was referenced only when needed to contextualize current standards.

Evidence was appraised qualitatively for relevance, methodological quality, and applicability to dental and head-and-neck settings. Where evidence originated in other surgical populations, such as abdominal surgery or general ward care, its transferability is stated explicitly, because extrapolation is a recurring limitation in this field. Findings were organized by professional domain and then synthesized into an integrated model. Because a narrative approach does not employ formal risk-of-bias scoring or quantitative synthesis, conclusions should be interpreted as informed expert integration rather than definitive effect estimates.

3. The Risk Landscape in Dental and Head-and-Neck Sedation and Anesthesia

3.1 Physiological and procedural hazards

Sedation exists on a continuum, and patients can progress unintentionally from minimal sedation to deep sedation or general anesthesia, losing protective reflexes and adequate ventilation. Contemporary practice guidelines emphasize that anyone administering sedation must be able to recognize and rescue a patient from a deeper level than intended, and that monitoring should be proportionate to the depth achieved (Homma et al., 2020). For pediatric patients, guidance stresses continuous observation by a person dedicated to monitoring, appropriate fasting and risk assessment, and readiness to manage airway compromise (Coté & Wilson, 2019).

Opioids and sedative-hypnotics depress ventilatory drive and upper airway tone, which is especially hazardous when the airway is shared with a surgical field. In a large prospective observational trial, continuous capnography and oximetry helped identify patients at risk of opioid-induced respiratory depression on inpatient wards, illustrating how physiological monitoring can reveal hypoventilation that intermittent checks miss (Khanna et al., 2020). Patients with obstructive sleep apnea represent a particular concern, and dedicated guidance recommends tailored preoperative identification, careful opioid use, and enhanced postoperative monitoring (Memtsoudis et al., 2018).

3.2 Airway complexity and the burden of preventable events

Analyses of closed claims have shown that difficult tracheal intubation continues to produce serious injury, and that unanticipated difficulty, repeated attempts, and delayed use of rescue strategies are common themes (Joffe et al., 2019). Updated national and international guidance stresses planning before induction, explicit communication of a primary and rescue strategy, and preparation for front-of-neck access when anticipated difficulty exists (Apfelbaum et al., 2022; Law et al., 2021). For patients with tumors, radiation fibrosis, or limited mouth opening, awake tracheal intubation is a core technique, and structured guidance exists for its safe performance (Ahmad et al., 2020). Such plans depend on information generated outside the anesthesia team, including imaging descriptions of airway anatomy and oncologic staging.

Capnography deserves special emphasis. Reviews of airway-related deaths argue that continuous waveform capnography would have prevented a considerable share of avoidable events by providing early recognition of obstruction, displacement, or apnea (Cook & Harrop-Griffiths, 2019). International monitoring standards now treat capnography as an essential component of care during anesthesia and recovery (Klein et al., 2021), and a Cochrane review found that adding capnography to standard monitoring during procedural sedation improved detection of respiratory depression in emergency settings (Wall et al., 2017). The global standards for safe anesthesia practice similarly define minimum requirements for personnel, equipment, and monitoring that apply irrespective of resource level (Gelb et al., 2018).

3.3 Developmental and perioperative considerations in children

Children undergoing dental rehabilitation represent a large share of sedation and anesthesia exposure in dentistry. Concerns about anesthetic neurotoxicity in the developing brain have prompted extensive investigation. Narrative synthesis suggests that brief, single exposures are unlikely to produce clinically meaningful deficits, although uncertainty remains regarding prolonged or repeated exposure (Andropoulos, 2018). A population-based cohort found no strong evidence that single exposures to anesthesia were associated with a decline in overall neuropsychological outcomes, while noting some differences in specific domains after multiple exposures (Warner et al., 2018). For safety planning, these findings support avoiding unnecessary or repeated anesthetics, consolidating treatment where clinically appropriate, and investing in prevention and behavioral management, a task that places dentistry at the center of risk reduction. Cochrane evidence on sedation for children undergoing dental treatment further highlights heterogeneity among agents and the need for standardized outcomes (Ashley et al., 2018).

3.4 Why interprofessional failure matters

The Swiss cheese model of accident causation remains a useful lens for these events. Harm occurs when weaknesses in several defensive layers align, whether these lie in preoperative assessment, equipment, supervision, communication, or organizational policy (Wiegmann et al., 2022). Teamwork science adds that high-performing teams share mental models, use closed-loop communication, monitor one another, and adapt to changing demands (Rosen et al., 2018). Figure 1 illustrates how each professional domain forms a protective layer around the patient and how coordination across layers reduces the likelihood of aligned failures.




Figure 1. Interprofessional layers of protection against preventable harm in dental and head-and-neck sedation and anesthesia (conceptual model adapted from the Swiss cheese perspective; Wiegmann et al., 2022).

4. Domain-Specific Contributions to Safety

4.1 Dentistry: risk prevention, patient selection, and procedural coordination

Dentists and oral and maxillofacial teams are often the first to assess the patient and therefore shape the entire risk trajectory. Several contributions are central. First, preventive care and behavioral guidance reduce the need for sedation or general anesthesia in the first place, which is the most effective way to eliminate anesthetic risk, particularly in children (Ashley et al., 2018; Andropoulos, 2018). Second, careful patient selection ensures that those with significant comorbidity, anticipated airway difficulty, or complex medication regimens are directed to settings with appropriate resources. Guidance on pediatric sedation stresses structured pre-sedation evaluation, informed consent, and clear criteria for the provider and facility (Coté & Wilson, 2019).

Third, dentists contribute to intraprocedural safety through the management of the shared airway. Packing, suction, isolation, and the use of throat protection reduce aspiration of debris, irrigants, or fragments, and the proceduralist must communicate clearly when intervention affects ventilation. For patients with obstructive sleep apnea or craniofacial abnormalities, dentists may identify anatomical risk markers and contribute to preoperative planning (Memtsoudis et al., 2018). Fourth, in head-and-neck oncology, dental teams manage pre-radiotherapy extractions and oral rehabilitation, and they follow patients for trismus, osteoradionecrosis, and infection, all of which influence later anesthetic access. Trismus, in particular, is a recognized sequela of head-and-neck cancer treatment that can restrict mouth opening and complicate laryngoscopy (Loh et al., 2017).

Finally, dental professionals must be competent in emergency response. Where sedation is provided in a dental office, immediate availability of oxygen, suction, airway adjuncts, monitoring, reversal agents, and personnel trained in basic and advanced life support is a prerequisite rather than an option, in line with global and national standards (Gelb et al., 2018; Klein et al., 2021). The interprofessional model suggests that the dentist's role is not merely to await anesthetic safety from the anesthesia provider but to co-own the pathway: selecting suitable patients, sharing airway information, and engaging in team briefings and debriefings.

4.2 Laboratory monitoring: risk stratification, bleeding, and metabolic safety

Laboratory services contribute to safety before, during, and after the procedure. The central principle in contemporary guidance is that testing should be targeted to clinical need rather than performed routinely. Cardiovascular guidelines for non-cardiac surgery encourage risk-based use of biomarkers and electrocardiography according to surgical risk and patient characteristics (Halvorsen et al., 2022). Over-testing generates false positives, delays, and cost without improving outcomes, while under-testing in patients with relevant comorbidity can miss modifiable risk.

Three laboratory domains are especially relevant. The first is hematological status. Anemia and iron deficiency are common in patients with head-and-neck cancer and are associated with worse perioperative outcomes, and consensus guidance recommends early detection and treatment as part of patient blood management (Muñoz et al., 2017). The second is hemostasis. Dental extractions and head-and-neck surgery can cause substantial bleeding in patients taking anticoagulants, antiplatelet agents, or with hepatic impairment, and bleeding into the airway can be rapidly life-threatening. European guidelines on severe perioperative bleeding support structured assessment, point-of-care viscoelastic testing where appropriate, and transfusion protocols driven by laboratory data (Kietaibl et al., 2023). The third domain is glycemic and metabolic control. Perioperative hyperglycemia is associated with infection and delayed healing, and updated recommendations support targeted screening, monitoring, and treatment pathways (Duggan et al., 2017).

The interprofessional dimension is the critical element. A laboratory result is only protective if it reaches the right person at the right time. Critical-value reporting, defined turnaround times for urgent testing, and integration of results into the preoperative checklist convert laboratory information into action. Laboratory scientists can also support point-of-care testing governance, ensuring that devices used in clinics are calibrated, quality-controlled, and interpreted correctly. Perioperative standards of complication definition, such as those developed for pulmonary and other endpoints, give laboratories, clinicians, and administrators a common vocabulary for auditing outcomes (Abbott et al., 2018). Table 1 summarizes the specific contributions of each professional group.

Table 1. Interprofessional contributions to safety in dental and head-and-neck sedation and anesthesia

Domain	Primary safety contribution	Key actions	Representative evidence
Dentistry	Prevention, patient selection, shared-airway management	Behavioral guidance; risk screening; isolation and suction; emergency competence; pre-radiotherapy dental care	Ashley et al. (2018); Coté & Wilson (2019); Loh et al. (2017)
Laboratory monitoring	Risk stratification and detection of metabolic or hemostatic risk	Targeted testing; anemia and glycemic optimization; coagulation and point-of-care testing; critical-value escalation	Kietaibl et al. (2023); Muñoz et al. (2017); Duggan et al. (2017)
Radiological technology	Anatomical risk mapping and procedural guidance	Airway and tumor imaging review; cone-beam CT; contrast and MRI safety; radiation stewardship; AI-assisted reporting	Kaasalainen et al. (2021); American College of Radiology (2015); Hosny et al. (2018)
Nursing	Continuous surveillance, rescue, and communication	Capnography and early-warning scores; sedation monitoring; PONV control; structured handover	Gerry et al. (2020); Gan et al. (2020); Müller et al. (2018)
Physiotherapy	Prevention of pulmonary, functional, and swallowing complications	Prehabilitation; breathing and mobilization; trismus, shoulder, and swallowing rehabilitation	Boden et al. (2018); Hutcheson et al. (2017); Loh et al. (2017)
Health administration	Systems, culture, staffing, and learning	Standards and credentialing; staffing; incident reporting; audits; simulation; equipment governance	WHO (2021); Braithwaite et al. (2017); Kruk et al. (2018)

4.3 Radiological technology: anatomical risk mapping and imaging safety

Radiological technology provides the anatomical map on which anesthetic planning depends. In head-and-neck procedures, computed tomography, magnetic resonance imaging, and ultrasound characterize tumor extent, airway narrowing, vascular relationships, and the presence of extrinsic compression. When such images are reviewed prospectively by the anesthesia team, they can inform the decision between asleep and awake intubation, the choice of airway device, and the need for surgical airway readiness, consistent with

recommendations that the plan for an anticipated difficult airway be formulated before induction (Apfelbaum et al., 2022; Law et al., 2021; Ahmad et al., 2020).

In dentistry, cone-beam computed tomography has become widely used for implant planning, impacted teeth, pathology, and craniofacial assessment. An updated review describes the technique's capabilities, the importance of justification and optimization, and the variability of radiation dose across devices and protocols (Kaasalainen et al., 2021). For sedated patients, this emphasizes that the radiographer's role includes dose stewardship, positioning that is safe for a sedated or anesthetized patient, and rapid acquisition to minimize time under sedation. Children, in particular, require attention to the principle of keeping exposure as low as reasonably achievable.

Sedation for imaging itself is a safety domain. Magnetic resonance environments constrain access to the patient and require MRI-conditional monitoring and equipment, so that radiographers, nurses, and anesthesia providers must share screening procedures and emergency plans. Where contrast media are used, institutional guidance addresses risk assessment for allergic-like reactions and renal considerations, premedication decisions, and treatment algorithms for adverse reactions (American College of Radiology, 2015). These processes depend on pre-procedure screening data supplied by laboratory services, such as renal function, which is an example of the cross-domain information flows that interprofessional models aim to secure.

Emerging digital tools may add further value. Artificial intelligence in radiology has demonstrated potential for automated segmentation, detection, and quantitative feature extraction that may support reporting efficiency and airway or tumor assessment (Hosny et al., 2018). Their safe use in perioperative workflows will require validation, transparency, and clear accountability, themes that fall within health administration. A final contribution lies in intraprocedural and postprocedural imaging, for instance to confirm positioning, detect complications such as aspiration or pneumothorax, or evaluate swelling. Radiographers who understand the clinical context can help prioritize urgent studies and communicate critical findings promptly, closing the loop between image acquisition and patient care.

4.4 Nursing: continuous surveillance, rescue, and communication

Nurses are the professional group with the most continuous contact with the sedated or recovering patient, and their observations frequently provide the earliest warning of deterioration. In sedation, the monitoring nurse or dedicated observer is responsible for assessing level of consciousness, respiratory pattern, oxygenation, ventilation, hemodynamics, and pain, and for escalating promptly when thresholds are crossed. Guidance on moderate and pediatric sedation requires a trained individual whose sole task is patient monitoring and who is able to initiate rescue (Homma et al., 2020; Coté & Wilson, 2019). Combined with continuous capnography, nursing observation provides a layered defense against unrecognized hypoventilation (Cook & Harrop-Griffiths, 2019; Klein et al., 2021).

Early-warning systems extend this surveillance into recovery and ward care. A systematic appraisal of early-warning scores found that, although many tools exist and several are widely implemented, methodological quality and validation vary considerably, which means that nurses and administrators should select tools suited to their population and verify performance locally (Gerry et al., 2020). Escalation pathways that empower nurses to call for senior or anesthetic help without fear of hierarchy are as important as the scoring tool itself, because a score that is not acted upon has no protective value. For patients at risk of opioid-induced respiratory depression, continuous monitoring with capnography and oximetry may be justified in selected groups (Khanna et al., 2020).

Nurses also lead on several complication-prevention bundles. Postoperative nausea and vomiting is a common cause of distress and, in head-and-neck and oral surgery, may increase the risk of bleeding, wound stress, and aspiration. Consensus guidelines recommend risk stratification and multimodal prophylaxis, tasks that nurses implement and audit at the bedside (Gan et al., 2020). Airway protection, positioning, oral care, suction, and tracheostomy or drain management are further nursing responsibilities with direct safety consequences, particularly in free-flap and oncologic head-and-neck surgery, where enhanced recovery recommendations emphasize standardized nursing care elements across the pathway (Dort et al., 2017).

Communication is the final and perhaps most underestimated nursing contribution. A systematic review of the SBAR handover tool found that structured communication was associated with improvements in patient safety outcomes and communication quality, although study designs were heterogeneous (Müller et al., 2018). In dental and head-and-neck pathways, handovers occur between the clinic, the imaging suite, the operating room, recovery, and the ward or home, and each transition is a point at which information about airway difficulty, laboratory abnormalities, sedation response, and rehabilitation needs can be lost. Finally, nurse staffing is itself a safety variable. A systematic review found that lower nurse staffing is associated with more omissions of necessary care, such as monitoring and mobilization (Griffiths et al., 2018), which links the bedside to the organizational decisions discussed in Section 4.6.

4.5 Physiotherapy: preventing pulmonary, functional, and swallowing complications

Although physiotherapists do not administer sedation, their work shapes how well patients tolerate it and how quickly they recover. A major contribution is the prevention of postoperative pulmonary complications. A pragmatic multicentre randomized trial of preoperative physiotherapy before upper abdominal surgery, combining education with breathing exercises, evaluated whether such preparation reduces postoperative pulmonary complications, and it remains one of the most direct trial investigations of this approach in a surgical population (Boden et al., 2018). Although this evidence derives from abdominal rather than head-and-neck surgery, the physiological rationale is relevant for patients with chronic lung disease, smoking history, or reduced reserve undergoing prolonged head-and-neck procedures, and it should be interpreted as plausible but not definitive transfer. Standardized definitions of pulmonary endpoints allow such interventions to be audited consistently (Abbott et al., 2018).

Prehabilitation and early mobilization are core components of enhanced recovery. Reviews of enhanced recovery after surgery describe multimodal pathways that include preoperative counseling and optimization, minimized fasting, opioid-sparing analgesia, and early mobilization, and report reductions in complications and length of stay across surgical specialties (Ljungqvist et al., 2017). Consensus recommendations specific to major head-and-neck surgery with free-flap reconstruction adapted these principles, giving a role to rehabilitation professionals in early mobilization, airway clearance, and function (Dort et al., 2017).

Head-and-neck physiotherapy and allied rehabilitation address specific functional sequelae. Trismus, which reduces mouth opening, can arise after surgery or radiotherapy and affects oral hygiene, eating, dental treatment, and airway access, and a systematic review has documented its prevalence and measurement across treatment modalities (Loh et al., 2017). Early jaw-stretching and mobility programs may therefore be considered both functional therapy and anesthetic risk-reduction, since improved mouth opening can facilitate future laryngoscopy. Shoulder dysfunction after neck dissection, cervical stiffness, and lymphedema are further areas where physiotherapists provide specialized care. Swallowing safety is another critical area, and dysphagia and aspiration are major drivers of pneumonia after head-and-neck treatment. Standardized instrumental grading systems, such as the dynamic imaging grade of swallowing toxicity, allow clinicians, radiographers, and speech-language therapists to describe swallowing physiology in a common language (Hutcheson et al., 2017). Physiotherapists collaborating with swallowing specialists and nurses contribute to chest physiotherapy, positioning, and respiratory training that lower the consequence of aspiration events.

In the interprofessional model, physiotherapy should be engaged before the procedure for risk assessment and prehabilitation, during the perioperative period through early mobilization, and after discharge through structured rehabilitation. When physiotherapy is introduced only after complications arise, its preventive potential is lost.

4.6 Health administration systems: governance, culture, and learning

Health administration systems determine whether the contributions of individual professions are reliable, reproducible, and sustained. The WHO global action plan identifies policy and leadership, safety culture, workforce capability, risk management, patient engagement, and learning systems as strategic pillars (WHO, 2021). In anesthesia specifically, international standards describe minimum requirements for facilities, equipment, monitoring, and personnel, and administrators are responsible for ensuring that these standards are met in every location where sedation is delivered, including dental clinics and imaging suites (Gelb et al., 2018). Because the global quality agenda stresses that poor-quality care now accounts for more deaths than lack of access in many low- and middle-income settings, safety investment is inseparable from health-system strengthening (Kruk et al., 2018).

Several administrative domains are particularly important. The first is credentialing and competency maintenance. Sedation and airway skills decay without practice, so recurring simulation, advanced life-support training, and supervised experience are required. Interprofessional education frameworks provide a structure for shared competencies in values and ethics, roles and responsibilities, communication, and teamwork, and they are intended to prepare professionals to work together before they meet at the bedside (Collaborative, 2023). The second domain is staffing and workload. Adequate nurse staffing and protected time for monitoring and handover are preconditions for surveillance (Griffiths et al., 2018).


The third domain is culture and organizational learning. A systematic review found that more positive organizational and workplace cultures, including leadership engagement and teamwork climate, are associated with better patient outcomes (Braithwaite et al., 2017). Incident reporting, structured debriefing, and open review of near misses transform individual events into system improvement, and the Swiss cheese perspective helps leaders look beyond individual blame to the latent conditions that allowed failure (Wiegmann et al., 2022). The fourth domain is information systems and equipment governance: electronic records that make laboratory values, imaging findings, and airway alerts visible at the point of care; maintenance schedules for monitors, capnographs, and difficult-airway carts; and standardized checklists for facilities that deliver sedation. Finally, economic analysis supports investment, since preventing harm is generally cost-

effective compared with treating complications (Slawomirski & Klazinga, 2022).

5. An Integrated Interprofessional Model

The preceding sections show that each profession contributes a necessary but insufficient layer of protection. Figure 2 proposes an integrated pathway that follows the patient through five phases: referral and screening, preoperative optimization and planning, the procedure, recovery, and rehabilitation and follow-up. At each phase, specific information flows between professions: dental risk assessment feeds laboratory and imaging requests; laboratory and imaging results inform the anesthesia and airway plan; nursing observations during and after the procedure trigger rescue; and rehabilitation outcomes feed back into future risk assessment. Health administration systems form the enabling environment, providing standards, staffing, data infrastructure, and learning loops.


Figure 2. Integrated interprofessional pathway showing the main contribution of each profession at each phase of care, supported by health administration systems.



Three design principles underpin the model. The first is shared situational awareness. The use of team briefings before the procedure, with explicit discussion of airway plan, bleeding risk, imaging findings, emergency roles, and equipment, supports shared mental models, a mechanism identified as central to high-performing healthcare teams (Rosen et al., 2018). The second is redundancy through diverse monitoring: clinical observation, pulse oximetry, capnography, and early-warning scores overlap so that no single failure leaves the patient unprotected (Klein et al., 2021; Gerry et al., 2020). The third is closed-loop information transfer, in which each critical data point, such as an abnormal coagulation result or a radiological airway finding, has a defined recipient, a time frame, and a confirmation of receipt. These principles are consistent with evidence that team processes are associated with performance and clinical outcomes (Schmutz et al., 2019) and that structured collaborative interventions can improve care when implemented well (Reeves et al., 2017).

The model also supports risk-tiered pathways. Low-risk patients undergoing minimal or moderate sedation in appropriate settings may follow streamlined pathways with limited testing and standard monitoring. Higher-risk patients, such as those with significant comorbidity, obstructive sleep apnea, anticipated difficult airway, prior head-and-neck radiotherapy, anticoagulation, or young age with extensive treatment needs, should be triaged to settings with anesthesia-led care, enhanced monitoring, and access to laboratory, imaging, and critical care support (Memtsoudis et al., 2018; Apfelbaum et al., 2022; Halvorsen et al., 2022). Figure 3 presents a continuous improvement cycle that administrators can use to sustain the model through measurement, audit, feedback, and revision.

Figure 3. Continuous improvement cycle for sustaining the interprofessional safety bundle through governance and audit.



6. Implementation Priorities

Translating the model into practice requires prioritization, because resources, personnel, and regulatory contexts differ. Table 2 outlines implementation priorities, responsible professions, and suggested indicators. The proposed indicators are not validated benchmarks; they are candidate measures that institutions can adapt, and their selection should follow local epidemiology and available data. Combining process indicators, such as compliance with capnography or briefing completion, with outcome indicators, such as unplanned airway intervention or pulmonary complications, helps distinguish adoption from effect.

Table 2. Implementation priorities, responsible professions, and candidate indicators

Priority	Lead and partner professions	Suggested actions	Candidate indicators
Standardized monitoring	Anesthesia, nursing, dentistry, administration	Continuous capnography and oximetry for all sedation; dedicated monitor; equipment checks	Percentage of cases with capnography; unplanned oxygen desaturation events

Priority	Lead and partner professions	Suggested actions	Candidate indicators
Risk-based preoperative pathway	Dentistry, laboratory, anesthesia	Targeted testing; anemia, glycemia, and anticoagulant plans; critical-value escalation	Cancellations due to missed results; time from critical value to action
Imaging-informed airway planning	Radiology, anesthesia, surgery	Prospective review of airway imaging; documented primary and rescue plan; awake intubation criteria	Percentage of high-risk cases with documented plan; unanticipated difficult airway rate
Structured communication	Nursing, all clinical teams	Pre-procedure briefing; SBAR handover; debriefing	Briefing compliance; handover-related incidents
Early rehabilitation	Physiotherapy, nursing, speech therapy	Prehabilitation; early mobilization; trismus and swallowing programs	Pulmonary complication rate; length of stay; mouth-opening change
Governance and learning	Administration, all professions	Credentialing; staffing standards; simulation; incident review; audit cycles	Staff completing simulation; incident reports reviewed; action closure rate

Successful implementation also depends on how change is introduced. Leadership commitment, local champions from several professions, protected training time, and feedback of audit data are repeatedly described as facilitators, while hierarchy, workload, and fragmented information systems act as barriers (Braithwaite et al., 2017; Bates & Singh, 2018). Interprofessional education delivered in simulation, with realistic scenarios such as a can't-intubate-can't-oxygenate event in a dental chair, provides a safe environment for rehearsing role clarity and communication (Collaborative, 2023). Equity considerations should also be explicit: low-resource clinics and rural facilities may lack capnography, laboratory access, or imaging, and global standards offer a framework for minimum safe provision and for phased improvement (Gelb et al., 2018; Kruk et al., 2018).

7. Research Gaps and Future Directions

Several gaps limit confident recommendations. First, high-quality interprofessional trials focused specifically on dental and head-and-neck sedation are scarce, and much of the supporting evidence is extrapolated from general surgical or inpatient populations (Boden et al., 2018; Khanna et al., 2020). Second, outcome definitions remain inconsistent; wider adoption of standardized perioperative endpoints would permit comparison across institutions and settings (Abbott et al., 2018). Third, the contribution of nursing staffing, physiotherapy, and laboratory turnaround to anesthetic outcomes in dental and head-and-neck pathways has rarely been measured directly. Fourth, evaluation of digital tools, including artificial intelligence-assisted imaging interpretation and electronic early-warning systems, requires prospective validation and attention to bias, interpretability, and workflow integration (Hosny et al., 2018; Gerry et al., 2020). Fifth, patient and family perspectives, particularly those of parents of children undergoing dental rehabilitation, are under-represented in safety research. Finally, implementation science studies are needed to identify which components of interprofessional bundles produce the greatest benefit per unit of resource, which is essential for settings with limited capacity.

8. Limitations

This review has limitations inherent in the narrative design. The search was not systematic, no formal risk-of-bias assessment or quantitative synthesis was performed, and selection of sources may reflect reviewer judgment. Much of the evidence derives from guidelines and consensus statements, whose recommendations may rest on indirect or low-certainty evidence. The restriction to 2017–2025 may have excluded earlier foundational studies, and the integrated model and candidate indicators are conceptual proposals that require empirical testing. Findings should be applied with attention to local regulation, resources, and patient populations.

9. Conclusion

Safety in sedation and anesthesia for dental and head-and-neck procedures is best understood as an emergent property of coordinated systems rather than the sum of individual competencies. Dentistry reduces exposure and shares ownership of the airway; laboratory monitoring detects hematological, hemostatic, and metabolic risk; radiological technology maps anatomy and protects patients during imaging; nursing provides continuous surveillance, rescue, and communication; physiotherapy prevents pulmonary, functional, and swallowing complications; and health administration provides the standards, staffing, culture, and learning loops that make these contributions reliable. The evidence reviewed supports standardized monitoring including capnography, targeted risk-based testing, imaging-informed airway planning, structured communication, early rehabilitation, and governance that supports competence and learning. Institutions should adopt these elements as an integrated package, measure both process and outcome, and share results so that the

evidence base specific to this high-risk interface continues to grow.

References

1. Abbott, T. E. F., Fowler, A. J., Pelosi, P., De Abreu, M. G., Møller, A. M., Canet, J., ... & Johnson, M. (2018). A systematic review and consensus definitions for standardised end-points in perioperative medicine: pulmonary complications. *British journal of anaesthesia*, *120*(5), 1066-1079.
2. Ahmad, I., El-Boghdadly, K., Bhagrath, R., Hodzovic, I., McNarry, A. F., Mir, F., ... & Vaughan, D. (2020). Difficult Airway Society guidelines for awake tracheal intubation (ATI) in adults. *Anaesthesia*, *75*(4), 509-528.
3. American College of Radiology. (2015). *ACR manual on contrast media*. American College of Radiology.
4. Andropoulos, D. B. (2018). Effect of anesthesia on the developing brain: infant and fetus. *Fetal diagnosis and therapy*, *43*(1), 1-11.
5. Apfelbaum, J. L., Hagberg, C. A., Connis, R. T., Abdelmalak, B. B., Agarkar, M., Dutton, R. P., ... & Tung, A. (2022). 2022 American Society of Anesthesiologists practice guidelines for management of the difficult airway. *Anesthesiology*, *136*(1), 31-81.
6. Ashley, P. F., Chaudhary, M., & Lourenço-Matharu, L. (2018). Sedation of children undergoing dental treatment. *Cochrane Database of Systematic Reviews*, (12).
7. Bates, D. W., & Singh, H. (2018). Two decades since to err is human: an assessment of progress and emerging priorities in patient safety. *Health Affairs*, *37*(11), 1736-1743.
8. Boden, I., Skinner, E. H., Browning, L., Reeve, J., Anderson, L., Hill, C., ... & Denehy, L. (2018). Preoperative physiotherapy for the prevention of respiratory complications after upper abdominal surgery: pragmatic, double blinded, multicentre randomised controlled trial. *bmj*, *360*.
9. Braithwaite, J., Herkes, J., Ludlow, K., Testa, L., & Lamprell, G. (2017). Association between organisational and workplace cultures, and patient outcomes: systematic review. *BMJ open*, *7*(11), e017708.
10. Collaborative, I. E. (2023). IPEC core competencies for interprofessional collaborative practice: Version 3. *Washington, DC: Interprofessional Education Collaborative*.
11. Cook, T. M., & Harrop-Griffiths, W. (2019). Capnography prevents avoidable deaths. *BMJ*, *364*.
12. Coté, C. J., Wilson, S., American Academy of Pediatrics, & American Academy of Pediatric Dentistry. (2019). Guidelines for monitoring and management of pediatric patients before, during, and after sedation for diagnostic and therapeutic procedures. *Pediatrics*, *143*(6), e20191000.
13. Dort, J. C., Farwell, D. G., Findlay, M., Huber, G. F., Kerr, P., Shea-Budgell, M. A., ... & Harris, J. (2017). Optimal perioperative care in major head and neck cancer surgery with free flap reconstruction: a consensus review and recommendations from the enhanced recovery after surgery society. *JAMA otolaryngology–head & neck surgery*, *143*(3), 292-303.
14. Duggan, E. W., Carlson, K., & Umpierrez, G. E. (2017). Perioperative hyperglycemia management: an update. *Anesthesiology*, *126*(3), 547.
15. Gan, T. J., Belani, K. G., Bergese, S., Chung, F., Diemunsch, P., Habib, A. S., ... & Philip, B. K. (2020). Fourth consensus guidelines for the management of postoperative nausea and vomiting. *Anesthesia & Analgesia*, *131*(2), 411-448.
16. Gelb, A. W., Morriss, W. W., Johnson, W., & Merry, A. F. (2018). World Health Organization-World Federation of Societies of Anaesthesiologists (WHO-WFSA) international standards for a safe practice of anesthesia. *Canadian Journal of Anesthesia/Journal canadien d'anesthésie*, *65*(6), 698-708.
17. Gerry, S., Bonnici, T., Birks, J., Kirtley, S., Virdee, P. S., Watkinson, P. J., & Collins, G. S. (2020). Early warning scores for detecting deterioration in adult hospital patients: systematic review and critical appraisal of methodology. *bmj*, *369*.
18. Griffiths, P., Recio-Saucedo, A., Dall'Ora, C., Briggs, J., Maruotti, A., Meredith, P., ... & Missed Care Study Group. (2018). The association between nurse staffing and omissions in nursing care: A systematic review. *Journal of advanced nursing*, *74*(7), 1474-1487.
19. Halvorsen, S., Mehilli, J., Cassese, S., Hall, T. S., Abdelhamid, M., Barbato, E., ... & Zacharowski, K. (2022). 2022 ESC Guidelines on cardiovascular assessment and management of patients undergoing non-cardiac surgery: Developed by the task force for cardiovascular assessment and management of patients undergoing non-cardiac surgery of the European Society of Cardiology (ESC) Endorsed by the European Society of Anaesthesiology and Intensive Care (ESAIC). *European heart journal*, *43*(39), 3826-3924.
20. Homma, Y., Norii, T., Kanazawa, T., Hoshino, A., Arino, S., Takase, H., ... & Japan

Society of Procedural Sedation and Analgesia. (2020). A mini-review of procedural sedation and analgesia in the emergency department. *Acute Medicine & Surgery*, *7*(1), e574.
21. Hosny, A., Parmar, C., Quackenbush, J., Schwartz, L. H., & Aerts, H. J. (2018). Artificial intelligence in radiology. *Nature Reviews Cancer*, *18*(8), 500-510.
22. Hutcheson, K. A., Barrow, M. P., Barringer, D. A., Knott, J. K., Lin, H. Y., Weber, R. S., ... & Lewin, J. S. (2017). Dynamic imaging grade of swallowing toxicity (DIGEST): scale development and validation. *Cancer*, *123*(1), 62-70.
23. Joffe, A. M., Aziz, M. F., Posner, K. L., Duggan, L. V., Mincer, S. L., & Domino, K. B. (2019). Management of difficult tracheal intubation: a closed claims analysis. *Anesthesiology*, *131*(4), 818.
24. Kaasalainen, T., Ekholm, M., Siiskonen, T., & Kortesniemi, M. (2021). Dental cone beam CT: An updated review. *Physica Medica*, *88*, 193-217.
25. Khanna, A. K., Bergese, S. D., Jungquist, C. R., Morimatsu, H., Uezono, S., Lee, S., ... & Overdyk, F. J. (2020). Prediction of opioid-induced respiratory depression on inpatient wards using continuous capnography and oximetry: an international prospective, observational trial. *Anesthesia and analgesia*, *131*(4), 1012.
26. Kietaibl, S., Ahmed, A., Afshari, A., Albaladejo, P., Aldecoa, C., Barauskas, G., ... & Zacharowski, K. (2023). Management of severe peri-operative bleeding: Guidelines from the European Society of Anaesthesiology and Intensive Care: Second update 2022. *European journal of anaesthesiology*, *40*(4), 226-304.
27. Klein, A. A., Meek, T., Allcock, E., Cook, T. M., Mincher, N., Morris, C., ... & Young, P. (2021). Recommendations for standards of monitoring during anaesthesia and recovery 2021: Guideline from the Association of Anaesthetists. *Anaesthesia*, *76*(9), 1212-1223.
28. Kruk, M. E., Gage, A. D., Arsenault, C., Jordan, K., Leslie, H. H., Roder-DeWan, S., ... & Pate, M. (2018). High-quality health systems in the Sustainable Development Goals era: time for a revolution. *The Lancet global health*, *6*(11), e1196-e1252.
29. Law, J. A., Duggan, L. V., Asselin, M., Baker, P., Crosby, E., Downey, A., ... & Canadian Airway Focus Group. (2021). Canadian Airway Focus Group updated consensus-based recommendations for management of the difficult airway: part 2. Planning and implementing safe management of the patient with an anticipated difficult airway. *Canadian Journal of Anesthesia/Journal canadien d'anesthésie*, *68*(9), 1405-1436.
30. Ljungqvist, O., Scott, M., & Fearon, K. C. (2017). Enhanced recovery after surgery: a review. *JAMA surgery*, *152*(3), 292-298.
31. Loh, S. Y., Mcleod, R. W., & Elhassan, H. A. (2017). Trismus following different treatment modalities for head and neck cancer: a systematic review of subjective measures. *European archives of oto-rhino-laryngology*, *274*(7), 2695-2707.
32. Memtsoudis, S. G., Cozowicz, C., Nagappa, M., Wong, J., Joshi, G. P., Wong, D. T., ... & Chung, F. (2018). Society of anesthesia and sleep medicine guideline on intraoperative management of adult patients with obstructive sleep apnea. *Anesthesia and analgesia*, *127*(4), 967.
33. Müller, M., Jürgens, J., Redaèlli, M., Klingberg, K., Hautz, W. E., & Stock, S. (2018). Impact of the communication and patient hand-off tool SBAR on patient safety: a systematic review. *BMJ open*, *8*(8), e022202.
34. Muñoz, M., Acheson, A. G., Auerbach, M., Besser, M., Habler, O., Kehlet, H., ... & Klein, A. A. (2017). International consensus statement on the peri-operative management of anaemia and iron deficiency. *Anaesthesia*, *72*(2), 233-247.
35. Panagioti, M., Khan, K., Keers, R. N., Abuzour, A., Phipps, D., Kontopantelis, E., ... & Ashcroft, D. M. (2019). Prevalence, severity, and nature of preventable patient harm across medical care settings: systematic review and meta-analysis. *bmj*, *366*.
36. Reeves, S., Pelone, F., Harrison, R., Goldman, J., Zwarenstein, M., & Cochrane Effective Practice and Organisation of Care Group. (1996). Interprofessional collaboration to improve professional practice and healthcare outcomes. *Cochrane database of systematic reviews*, *2018*(8).
37. Rosen, M. A., DiazGranados, D., Dietz, A. S., Benishek, L. E., Thompson, D., Pronovost, P. J., & Weaver, S. J. (2018). Teamwork in healthcare: Key discoveries enabling safer, high-quality care. *American Psychologist*, *73*(4), 433.
38. Schmutz, J. B., Meier, L. L., & Manser, T. (2019). How effective is teamwork really? The relationship between teamwork and performance in healthcare teams: a systematic review and meta-analysis. *BMJ open*, *9*(9), e028280.
39. Slawomirski, L., & Klazinga, N. (2022). The economics of patient safety: from analysis to action. *OECD Health Working Papers*.

40. Wall, B. F., Magee, K., Campbell, S. G., & Zed, P. J. (2017). Capnography versus standard monitoring for emergency department procedural sedation and analgesia. *Cochrane Database of Systematic Reviews*, (3).
41. Warner, D. O., Zaccariello, M. J., Katusic, S. K., Schroeder, D. R., Hanson, A. C., Schulte, P. J., ... & Flick, R. P. (2018). Neuropsychological and behavioral outcomes after exposure of young children to procedures requiring general anesthesia: The MASK study. *Anesthesiology*, *129*(1), 89.
42. Wiegmann, D. A., Wood, L. J., Cohen, T. N., & Shappell, S. A. (2022). Understanding the "Swiss cheese model" and its application to patient safety. *Journal of patient safety*, *18*(2), 119.

World Health Organization. (2021). *Global patient safety action plan 2021-2030: towards eliminating avoidable harm in health care*. World Health Organization.